


# VITACK<sub>3</sub>RI



VITACK<sub>3</sub>RI is used in all animal species, including volatiles, to meet the need for vitamins and minerals.

#### USAGE AREA:

VITACK<sub>3</sub>RI is used for Vitamin A, C, and K deficiencies, growth, and development in newborn and young animals used to increase body resistance against bacterial diseases and used to support the organism. It is used for the treatment of poisoning (especially in mycotoxicosis in volatiles). It is used to eliminate the negative effects of stress factors related to the environment, habitat, and climate change. It is used for treatment in various bleeding cases in the skin and muscles of volatiles. It is used to assist in the reduction of reproduction and yield caused by the deficiency or absence of vitamins in the formulation.

#### USAGE:

It is used in the following way for 3-5 days by adding to the drinking water:

It is given to volatiles for 3-5 days by adding 100 g to 200 Lt (1 barrel) of drinking water.

It is given to other animals for 3-5 days by adding to the drinking water and mixing in water with the calculation of 5 g (1 scale) per 50 kg live weight.

The mixture should be prepared fresh every day and consumed within 2-3 hours.

#### METHOD OF COMMERCIAL PRESENTATION:

It is offered for sale in 100 Gr doypack packages and 1 Kg plastic jars.

# ViTACK<sub>3</sub>RI

ViTACK <sub>3</sub> RI				
Active Ingredient	Contribution Name	Premix Level (Every 25 KG)	Units	Identify Number
<b>Vitamins, Provitamins and Well-Identfy Chemicals with Similar Biological Effects to Vitamins</b>				
Vitamin A (Beta Carotene)	Vitamin A 500.000 IU	4.000.000	IU	E672
Vitamin C (Ascorbic Acid)	Vitamin C %99	10.000	mg	E300
Vitamin K <sub>3</sub>	Vitamin K <sub>3</sub> %51	3.000	mg	
<b>Minerals and Their Products</b>				
Sodium Bicarbonate	Sodium Bicarbonate	50.000	mg	11.4.2
<b>Other Beits</b>				
Maltodextrin	Maltodextrin	QSP	mg	13.3.6

